

Micro-domains as site for storage lipid biosynthesis in seeds and algae

Dissertation

for the award of the degree

“Doctor rerum naturalium (Dr. rer. nat.)”

of the Georg-August-University Göttingen

within the doctoral program “GGNB Microbiology and Biochemistry”

of the Georg-August-University School of Science (GAUSS)

submitted by

Daniel Bruckhoff

from Emmerich am Rhein, Germany

Göttingen, 2014

Bibliografische Information der Deutschen Bibliothek
Die Deutsche Bibliothek verzeichnet diese Publikation in der Deutschen Nationalbibliografie; detaillierte bibliografische Daten sind im Internet über <http://dnb.ddb.de> abrufbar.

Members of the Thesis Committee:

Prof. Dr. Ivo Feußner
Department for Plant Biochemistry, Albrecht-von-Haller-Institute for Plant Sciences,
University of Göttingen
Prof. Dr. Jörg Stülke
Department of General Microbiology, University of Göttingen
Prof. Dr. Andrea Polle
Department for Forest Botany and Tree Physiology, University of Göttingen

Members of the Examination Board:

Reviewer Prof. Dr. Ivo Feußner
Department for Plant Biochemistry, Albrecht-von-Haller-Institute for Plant Sciences,
University of Göttingen
Reviewer Prof. Dr. Jörg Stülke
Department of General Microbiology, University of Göttingen
Prof. Dr. Christiane Gatz
Department of Plant Molecular Biology and Physiology, Schwann-Schleiden-Research
Center for Molecular Cell Biology, University of Göttingen
PD Dr. Franz Hadacek
Department for Plant Biochemistry, Albrecht-von-Haller-Institute for Plant Sciences,
University of Göttingen
Prof. Dr. Andrea Polle
Department for Forest Botany and Tree Physiology, University of Göttingen
PD Dr. Thomas Teichmann
Department of Plant Cell Biology, Schwann-Schleiden-Research Center
for Molecular Cell Biology, University of Göttingen
Date of the oral examination: January 5, 2014

Bruckhoff, Daniel:

Micro-domains as site for storage lipid biosynthesis in seeds and algae
ISBN 978-3-86376-171-4

Alle Rechte vorbehalten
1. Auflage 2015
© Optimus Verlag, Göttingen
© Coverfoto: Daniel Bruckhoff
Coverlayout: Dipl.-Kfm. Alexander Mostafa
URL: www.optimus-verlag.de, Printed in Germany,
Papier ist FSC zertifiziert (holzfrei, chlorfrei und säurefrei
sowie alterungsbeständig nach ANSI 3948 und ISO 9706)

Das Werk, einschließlich aller seiner Teile, ist urheberrechtlich geschützt. Jede Verwertung außerhalb der engen Grenzen des Urheberrechtsgesetzes in Deutschland ist ohne Zustimmung des Verlages unzulässig und strafbar. Dies gilt insbesondere für Vervielfältigungen, Übersetzungen, Mikroverfilmungen und die Einspeicherung und Verarbeitung in elektronischen Systemen.

Table of Contents

List of Figures	V
List of Tables	VIII
Abbreviations	IX
Acknowledgements.....	XI
1 Introduction	1
1.1 Increasing need for triacylglycerol.....	1
1.2 Triacylglycerol (TAG)	4
1.2.1 TAG synthesis	6
1.3 Lipid droplets (LDs)	10
1.4 Oleosins	12
1.5 Diacylglycerol acyltransferases	13
1.6 Arabidopsis acyl-CoA binding protein 6 (AtACBP6)	14
1.7 Aim of the thesis.....	15
2 Material and methods	17
2.1 Material.....	17
2.1.1 Technical equipment	17
2.1.2 Chemicals and standards.....	18
2.1.3 Antibiotics	18
2.1.4 Consumables	19
2.1.5 Enzymes, Polymerases and size markers	19
2.1.6 Kits.....	20
2.1.7 Plasmids.....	20
2.1.8 Oligonucleotides for DNA constructs.....	21
2.1.9 Software	22
2.1.10 Bacteria	23

2.1.11 <i>S. cerevisiae</i>	23
2.1.12 Plants	24
2.1.13 Algae	24
2.2 Methods	24
2.2.1 Molecular biology	24
2.2.1.1 Polymerase chain reaction (PCR)	24
2.2.1.2 Standard PCR	25
2.2.1.3 Touch down PCR	26
2.2.1.4 Fusion of genes by PCR (fusion PCR)	26
2.2.1.5 Colony PCR	27
2.2.1.6 Semi-quantitative RT-PCR	28
2.2.1.7 Horizontal gel electrophoresis	28
2.2.1.8 Sequencing	29
2.2.1.9 Subcloning	29
2.2.1.10 Restriction and Ligation	29
2.2.2 Maintenance and treatment of <i>E. coli</i>	29
2.2.2.1 Growth conditions of <i>E. coli</i>	30
2.2.2.2 Preparation of chemically competent <i>E. coli</i> cells	30
2.2.2.3 Transformation of <i>E. coli</i> cells	30
2.2.3 Maintenance and treatment of <i>A. tumefaciens</i>	31
2.2.3.1 Growth conditions of <i>A. tumefaciens</i>	31
2.2.3.2 Preparation of chemically competent <i>A. tumefaciens</i> cells	31
2.2.3.3 Transformation of <i>A. tumefaciens</i> cells	32
2.2.4 Maintenance and treatment of <i>S. cerevisiae</i>	32
2.2.4.1 Growth conditions of <i>S. cerevisiae</i>	32
2.2.4.2 Preparation of competent <i>S. cerevisiae</i> cells	32
2.2.4.3 Transformation of <i>S. cerevisiae</i> cells	33
2.2.4.4 Feeding with FA or fatty alcohols	33
2.2.5 Maintenance and treatment of Arabidopsis plants	34
2.2.5.1 Growth conditions of Arabidopsis	34

2.2.5.2	Transformation of Arabidopsis plants	34
2.2.5.3	Greenhouse experiments	35
2.2.6	Maintenance and treatment of <i>P. tricornutum</i>	36
2.2.6.1	Growth conditions	36
2.2.6.1	Transformation control of <i>P. tricornutum</i>	38
2.2.7	Isolation of DNA and RNA	38
2.2.7.1	Isolation of genomic DNA from Arabidopsis and <i>P. tricornutum</i>	38
2.2.7.2	Isolation of plasmid DNA	39
2.2.7.3	RNA isolation by TRIzol	39
2.2.7.4	RNA isolation from <i>S. cerevisiae</i>	40
2.2.7.5	RNA isolation from siliques	41
2.2.7.6	cDNA synthesis	42
2.2.8	Biochemical methods	42
2.2.8.1	Lipid analysis from total lipid extracts from <i>S. cerevisiae</i>	42
2.2.8.2	Lipid extraction from <i>P. tricornutum</i>	43
2.2.8.3	Lipid extraction from Arabidopsis seeds and acidic methanolysis	44
2.2.8.4	DGAT functionality assay	44
2.2.8.5	Gas-chromatography with flame ionization detection (GC-FID)	44
3	Results	47
3.1	Heterologous expression in <i>S. cerevisiae</i>	47
3.1.1	Heterologous expression of EhDGAT2 in <i>S. cerevisiae</i>	48
3.1.1.1	Feeding with fatty acids and fatty alcohols	50
3.1.2	Homologous expression of AtOLEO3, AtDGAT1 and AtOLEO3-AtDGAT1 in <i>S. cerevisiae</i>	52
3.1.3	Heterologous expression of AtDGAT3	53
3.1.3.1	Feeding with fatty acids	54
3.2	Homologous expression in <i>Arabidopsis thaliana</i>	55
3.2.1	FA analysis of T3 seeds of construct 3 (napin::AtDGAT1)	57

3.2.2 FA analysis of T3 seeds of construct 64 (napin::AtACBP6, napin::AtDGAT1).....	60
3.2.3 FA analysis of T3 seeds of construct 65 (napin::AtOLEO3, napin::AtDGAT1)	63
3.2.4 FA analysis of T3 seeds of construct 74 (napin::AtOLEO3-AtDGAT1)	66
3.3 Heterologous expression in <i>P. tricornutum</i>	71
3.3.1 Transformation control by PCR	71
3.3.2 Growth kinetic of <i>P. tricornutum</i>	73
3.3.3 TAG analysis of <i>P. tricornutum</i>	75
4 Discussion	83
4.1 Heterologous expression in <i>S. cerevisiae</i>	83
4.1.1 Heterologous expression of EhDGAT2 in <i>S. cerevisiae</i>	83
4.1.2 Heterologous expression of AtOLEO3, AtDGAT1 and AtOLEO3-AtDGAT1 in <i>S. cerevisiae</i>	86
4.1.3 Heterologous expression of AtDGAT3 in <i>S. cerevisiae</i>	87
4.2 Attempts to increase oil content in Arabidopsis seeds.....	91
4.2.1 FA analysis of seeds of construct 3 (napin::AtDGAT1).....	91
4.2.2 FA analysis of seeds of construct 64 (napin::AtACBP6, napin::AtDGAT1).....	93
4.2.3 FA analysis of seeds of construct 65 (napin::AtOLEO3, napin::AtDGAT1)	94
4.2.4 FA analysis of seeds of construct 74 (napin::AtOLEO3-AtDGAT1)	96
4.3 Heterologous expression in <i>P. tricornutum</i>	98
Summary	101
Appendix	103
Literature	149

List of Figures

Figure 1-1:	Centers of domestication.	2
Figure 1-2:	TAG molecule.	5
Figure 1-3:	Simplified model of TAG biosynthesis in plants.	7
Figure 1-4:	Different origins TAG.	10
Figure 1-5:	Lipid droplet (LD) formation.	11
Figure 1-6:	Targeting of ACBP6 and DGAT via AtOLEO3 fusions to LDs.	16
Figure 2-1:	Principle steps of a fusion PCR. Start codon of gene 1 and stop codon of gene 2 are substituted by a linker sequence by PCR. To fuse both genes a further PCR is performed using both modified genes as a template. The black arrows indicate PCRs.	27
Figure 2-2:	Arrangement of trays for a greenhouse experiment.	35
Figure 2-3:	Arrangement of plants on a tray.	36
Figure 3-1:	Gene expression of EhDGAT2 in the <i>S. cerevisiae</i> double mutant $\Delta dga1\Delta lro1$	49
Figure 3-2:	TLC of the quadruple mutant H1246 with EhDGAT2 and feeding with fatty acids and fatty alcohols.	50
Figure 3-3:	Complementation of the quadruple mutant H1246 complemented with AtOLEO3, AtDGAT1 and AtOLEO3-AtDAGAT1.	53
Figure 3-4:	TLC of the quadruple mutant H1246 transformed with AtDGAT3 and AtOLEO3-AtDGAT3 and feeding with fatty acids.	54
Figure 3-5:	Overview of the mean values for the total fatty acid content of all measured lines of construct 3 (napin::AtDGAT1) with corresponding EVCs and wild types.	57
Figure 3-6:	Expression analysis of construct 3 (napin::DGAT1) by semi-quantitative RT-PCR.	58

Figure 3-7:	Relative fatty acid composition of construct 3 (napin::AtDGAT1).	59
Figure 3-8:	Overview of the mean values for the total fatty acid content of all measured lines of construct 64 (napin::OLEO3, napin::AtDGAT1) with corresponding EVCs and wild types.....	60
Figure 3-9:	Expression analysis of construct 64 (napin::AtACBP6, napin::DGAT1) by semi-quantitative RT-PCR.	61
Figure 3-10:	Relative fatty acid composition of construct 64 (napin::AtOLEO3, napin::AtDGAT1).	62
Figure 3-11:	Overview of the mean values for the total fatty acid content of all measured lines of construct 65 (napin::OLEO3, napin::AtDGAT1) with corresponding EVCs and wild types.....	63
Figure 3-12:	Expression analysis of construct 65 (napin::AtOLEO3, napin::AtDGAT1) by semi-quantitative RT-PCR.....	64
Figure 3-13:	Relative fatty acid composition of construct 65 (napin::AtOLEO3, napin::AtDGAT1).	65
Figure 3-14:	Overview of the mean values for the lipid content of all measured lines of construct 74 (napin::OLEO3-AtDGAT1) with corresponding EVCs and WT.	66
Figure 3-15:	Expression analysis of construct 74 (napin::AtOLEO3-AtDGAT1) by semi-quantitative RT-PCR.	67
Figure 3-16:	Expression analysis for the gene fusion AtOLEO3-AtDGAT1 in transformants of construct 76 (napin::ACBP6-AtOLEO3, napin::AtOLEO3-AtDGAT1) by semi-quantitative RT-PCR.	68
Figure 3-17:	Transformation analysis by PCR.	69
Figure 3-18:	Relative fatty acid composition of construct 74 (napin::AtOLEO3-AtDGAT1).....	70
Figure 3-19:	Transformation analysis of the construct Pt7 (fcpA::AtOLEO3(co)).....	71

Figure 3-20: Transformation analysis of the construct Pt16 (fcpA::AtDGAT1(co)).	72
Figure 3-21: Colony-PCR of the construct Pt18 (fcpA::AtOLEO3-AtDGAT1(co)).	72
Figure 3-22: Cell number of <i>P. tricornutum</i> cultures.	74
Figure 3-23: Chlorophyll a content of <i>P. tricornutum</i> cultures.	75
Figure 3-24: Total FA content of the <i>P. tricornutum</i> cultures.	76
Figure 3-25: TAG content of the <i>P. tricornutum</i> cultures.	78
Figure 3-26: FA content of the main FA of chain length C16 of the <i>P. tricornutum</i> cultures.	79
Figure 3-27: Content of the main FA of chain length C20 and C22 of the <i>P. tricornutum</i> cultures.	80
Figure 3-28: Main FA with chain length C16 of the TAG fraction of <i>P. tricornutum</i> cultures.	82
Figure 4-1: Membrane prediction of EhDGAT2.	85
Figure 4-2: Alignment of DGAT3 from <i>Arachis hypogaea</i> (AhDGA3) and the two versions of from <i>Arabidopsis</i> AtDGAT3.	89
Figure 4-3: Alignment of AtACBP6 and BnACBP. The alignment was performed using an internet tool (http://www.ebi.ac.uk/Tools/msa/clustalw2/). Symbols: identical (asterisk), conserved (colon), semi-conserved (dot).	94

List of Tables

Table 2-1:	Technical equipment used during this thesis.....	17
Table 2-2:	Chemicals and standards used during this thesis.	18
Table 2-3:	Consumables used during this thesis.	19
Table 2-4:	Enzymes, Polymerases and size markers used during this thesis.	19
Table 2-5:	Kits used in this thesis.	20
Table 2-6:	Plasmids used in this thesis.....	20
Table 2-7:	Oligonucleotides used in this thesis. Sequences belonging to the gene are given in capital letters, restriction sites are underlined small letters that are not underlined are linker sequences.....	21
Table 2-8:	Software and web-based tools used in this thesis.....	22
Table 2-9:	Bacterial strains used in this thesis.....	23
Table 2-10:	<i>S. cerevisiae</i> strains used in this thesis.	23
Table 2-11:	Constructs transformed into Arabidopsis.	24
Table 2-12:	Temperature program of a standard PCR.	25
Table 3-1:	Constructs grown in green house for experiment 1 and experiment 2.	56
Table 4-1:	Comparison of the codon usage of <i>S. cerevisiae</i> for codon-optimization of EhDGAT2.	86
Table 4-2:	Comparison of the codon usage of <i>S. cerevisiae</i> for codon-optimization of AtDGAT3. Differences between the codon usage from the publication (Bennetzen et al 1981) and the codon usage from the internet tool (gcu.schoedl.de) are marked in bold letters.....	91

Abbreviations

A ₆₅₀	Absorbance at wavelength 650 nm of light
bp	base pairs
C	Cytosine
CTAB	Cetyltrimethylammonium bromide
cDNA	Complementary DNA
d	Day
DAG	Diacylglycerol
DMSO	Dimethyl sulfoxide
DNA	Deoxyribonucleic acid
dNTP	Dinucleotide triphosphate
DW	Dry weight
EDTA	Ethylenediaminetetra acetic acid
ER	Endoplasmic reticulum
<i>et al.</i>	<i>et alii</i>
FA	Fatty acid
FID	Flame ionization detector
G	Gramm
GC	Gas chromatograph
h	Hours
H ₂ O	Water
l	Liters
LB	Luria bertani
m	Meters
M	Mol
MetOH	Methanol
min	Minutes
OD ₆₀₀	Optical density at 600 nm wavelength of light
PCR	Polymerase chain reaction
RNA	Ribonucleic acid
rpm	Rounds per minute
RT	Reverse transcriptase
s	Seconds
SDS	Sodium dodecyl sulphate
T	Thymine
TAG	Triacylglycerols
TLC	Thin layer chromatography
Tris	2-amino-2-hydroxymethyl-1,3-propanediol
URA 3	Selection marker for Uracil auxotrophy
v/v	Volume to Volume ratio
w/v	Weight to volume ratio
x g	Gravitation